



ABN 57 080 791 420

Integrated Nutrition Management Pty. Ltd.

38 Flaxton Mill Rd. Flaxton Qld. 4560 AUSTRALIA

Ph. (02) 8117 8154, Fax. (02) 8117 8155. Mobile 0417 408 795

Email: info@inmpl.com

LIQUID K-MAX

Nitrogen (0%), Phosphorus (8%), Potassium (25%)

Contains Fulvic Acids (0.5%)

Acid Buffered

LIQUID K-MAX is a liquid version of **K-MAX**. It is recommended for use by growers who prefer liquid products and/or in situations where water quality does not allow the use of solid products.

PRODUCT INFORMATION:

- **LIQUID K-MAX** is a foliar, totally soluble high potency Potassium and Phosphorus fertilizer.
- It is Nitrogen free.
- It contains Phosphorus at a ratio similar to that of Potassium in most crops.
- It contains soluble Fulvic Acids for better absorption and translocation.
- It is Acid Buffered. It will not increase the pH of juice sap of crops.
- It is water free, and whence extremely price effective.
- Its formulation guarantees immediate uptake and translocation in the plant tissue.

PROBLEMS IN DEALING WITH POTASSIUM DEFICIENCIES:

- They are widespread on most soils at times of stress.
- They are highly susceptible to nutritional imbalances.
- Potassium, in some instances, is locked up in the soil and unavailable to plants.
- Soil applications at times of stress are normally too slow to generate the right response quickly enough.

PROBLEMS WITH FOLIAR POTASSIUM FERTILIZERS:

- Some of them are salts of Sulphate, Nitrates and/or Chlorides. All of these forms are inefficient and lead to little response.
- They have a very low rate of translocation inside the plant.
- Most, if not all, foliar Potassium are liquids. Their end-user price includes the cost of packaging and freight.
- Most foliar Potassium products have a tendency to increase the pH of fruit juice.

DIRECTIONS FOR USE

APPLES:

2-3 Lt/Ha (depending on canopy size) when fruit reaches approx. 3rd full size.
2-3 Lt/Ha (depending on canopy size) approx. 4-6 weeks later.

CITRUS:

3-4 Lt/Ha (depending on canopy size) in later winter.
3-4 Lt/Ha (depending on canopy size) when fruit reaches approx. half size.

COTTON:

2-3 Lt/Ha at flowering.
2-3 Lt/Ha during early boll formation.

CUT FLOWERS:

2-3 Lt/Ha during early stem formation.
2-3 Lt/Ha just before bud formation.

FIELD CROPS:

1.5-2 Lt/Ha approx. 2 weeks after germination.
2-3 Lt/Ha approx. 2-3 weeks later.

GRAPES: (Table and Wine varieties)

2-3 Lt/Ha (depending on canopy size) at bunch closure.
2-3 Lt/Ha (depending on canopy size) approx. 3-4 weeks before picking.

GREENHOUSE CROPS:

500 ml/200 Lt water. Repeat application every 2-3 weeks

POTATO:

1.5-2 Lt/Ha at early tuber formation.
2-3 Lt/Ha soon after flowering.

STONE FRUIT:

1.5-2 Lt/Ha (depending on canopy) when fruit reaches half size.
1.5-2 Lt/Ha (depending on canopy size) 2-3 weeks before picking.

COMPATIBILITY:

LIQUID K-MAX is compatible with most agricultural chemical and fertilizers. Incompatible with Calcium. It is advised that a small compatibility test be done before mixing with other farm chemicals.

