



ABN 57 080 791 420

Integrated Nutrition Management Pty. Ltd.

38 Flaxton Mill Rd. Flaxton Qld. 4560 AUSTRALIA

Ph. (02) 8117 8154, Fax. (02) 8117 8155. Mobile 0417 408 795

Email: info@inmpl.com

MAG-MAX

Magnesium (Mg) 10%

Prevents and treats Magnesium deficiencies

PRODUCT INFORMATION:

- **MAG-MAX** is a **foliar** Magnesium product.
- **MAG-MAX** is specially formulated using organic chelating and complexing agents in order to assure better and quicker uptake and translocation within the plant tissue.
- **MAG-MAX** provides quick, efficient, safe, and **cost-effective** way of treating serious Magnesium deficiency disorders in most crops.
- **MAG-MAX** has a formulation that facilitates **efficient and prompt uptake** as well as translocation in the plant tissue.

MAGNESIUM DEFICIENCIES:

- Are widespread.
- Are very common, especially on acid soils.
- Unlike Trace Elements, Magnesium is a **Major Element**. To correct its deficiencies by foliar applications, many applications are required at high rates of application. It is always advisable to apply soil magnesium when magnesium is deficient.
- May be the result of **other nutritional imbalances**, especially high levels of Nitrogen, Calcium, & Phosphorus.

DIRECTIONS FOR USE

APPLES:

600 – 900 gr/Ha (depending on canopy size) 2-3 weeks after fruit set.

600 - 900 gr/Ha (depending on canopy size) when fruit reaches approx. half full size.

CITRUS:

800-1200 gr/Ha (depending on canopy size) when fruit reaches approx. half size.

800-1200 gr/Ha (depending on canopy size) as fruit colour turns orange

CUT FLOWERS:

700 gr/Ha during early stem formation.

800 gr/Ha just before bud formation.

GREENHOUSE CROPS:

300 gr/200 Lt water. Repeat application every 2-3 weeks

FIELD CROPS:

600 gr/Ha at 6-7 leaf stage.

800 gr/Ha 3-4 weeks later.

STONE FRUIT:

600-900 gr/Ha (depending on canopy) soon after petal fall.

600-900 gr/Ha (depending on canopy size) when fruit reaches half size.

600-900 gr/Ha (depending on canopy size) approx. 10-14 days before picking.

STRAWBERRY:

400-500 gr/Ha early in the season.

Repeat application every 3-4 weeks or as needed.

TOMATO:

700 gr/Ha at first cluster set.

Repeat application every 2-3 weeks.

FOR OTHER CROPS, APPLY 600-800 GR/HA (TWICE) OR SEEK THE ADVICE OF YOUR DISTRIBUTOR.

COMPATIBILITY:

MAG-MAX is compatible with most agricultural chemicals and fertilizers. It is however advised that a small compatibility test be done before mixing with other farm chemicals.

PACKAGING: 5 kg reusable plastic pails.

Sincerely

(electronic mail)

Iman SAFI/ Director